


This article was downloaded by: [VUL Vanderbilt University]

On: 18 April 2014, At: 07:28

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Cognitive Neuropsychiatry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/pcnp20>

Failure to benefit from target novelty during encoding contributes to working memory deficits in schizophrenia

Jutta S. Mayer^{ab}, Jejoong Kim^c & Sohee Park^a

^a Department of Psychology, Vanderbilt University, 111 21st Ave South, Nashville, TN 37240, USA

^b Department of Psychology, Goethe University, Postfach 11 19 32, Fach 128, D-60054 Frankfurt, Germany

^c Department of Psychology, Duksung Women's University, 419 Ssangmun-Dong, Dobong-Gu, Seoul 132-714, Korea

Published online: 11 Nov 2013.

To cite this article: Jutta S. Mayer, Jejoong Kim & Sohee Park (2014) Failure to benefit from target novelty during encoding contributes to working memory deficits in schizophrenia, *Cognitive Neuropsychiatry*, 19:3, 268-279, DOI: [10.1080/13546805.2013.854199](https://doi.org/10.1080/13546805.2013.854199)

To link to this article: <http://dx.doi.org/10.1080/13546805.2013.854199>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Failure to benefit from target novelty during encoding contributes to working memory deficits in schizophrenia

Jutta S. Mayer^{a,b}, Jejoong Kim^c and Sohee Park^{a*}

^aDepartment of Psychology, Vanderbilt University, 111 21st Ave South, Nashville, TN 37240, USA;

^bDepartment of Psychology, Goethe University, Postfach 11 19 32, Fach 128, D-60054 Frankfurt, Germany; ^cDepartment of Psychology, Duksung Women's University, 419 Ssangmun-Dong, Dobong-Gu, Seoul 132-714, Korea

(Received 4 March 2013; accepted 25 September 2013)

Introduction. Although working memory (WM) impairments are well documented in schizophrenic patients (PSZ), the underlying mechanisms are poorly understood. The aim of this study was to investigate the role of target salience during encoding to determine whether impaired visual attention in PSZ leads to poor WM.

Methods. Thirty-one PSZ and 28 demographically matched healthy controls (HC) performed a spatial delayed-response task. Attentional demands were manipulated during WM encoding by presenting high salient (novel) or low salient (familiar) targets. Participants also rated their level of response confidence at the end of each trial, allowing us to analyse different response types.

Results. WM was impaired in PSZ. Increasing target salience by increasing novelty improved WM performance in HC but not in PSZ. Poor WM performance in PSZ was largely due to an increase in the proportion of incorrect but high confident responses most likely reflecting a failure to encode the correct target.

Conclusions. Our findings suggest that dysfunctions of non-mnemonic attentional processes during encoding contribute to WM impairments in schizophrenia and may represent an important target for cognitive remediation strategies.

Keywords: spatial working memory; encoding; attention; schizophrenia; relatives; false memory

Impairments in working memory (WM) are core cognitive deficits in schizophrenia (Goldman-Rakic, 1994; Lee & Park, 2005) with significant consequences on social and occupational functioning (Cervellione, Burdick, Cottone, Rhinewine, & Kumra, 2007). Spatial WM deficits are present in high-risk populations (Smith, Park, & Cornblatt, 2006), in spectrum disorders (Mitropoulou et al., 2005), and in unaffected relatives (Myles-Worsley & Park, 2002; Park, Holzman, & Goldman-Rakic, 1995; Snitz, MacDonald, & Carter, 2006), and therefore have been proposed to be a potential endophenotypic marker of schizophrenia (Glahn, Therman, & Manninen, 2003; Meyer-Lindenberg & Weinberger, 2006; Saperstein et al., 2006). Although WM deficits in schizophrenic patients (PSZ) are well documented, it is still unclear what causes these deficits and whether they can be ameliorated. WM is a complex system of various subprocesses (Bledowski, Kaiser, & Rahm, 2010). For a clear understanding of the nature of WM impairments and to develop targeted remediation strategies, it is therefore crucial

*Corresponding author. Email: sohee.park@vanderbilt.edu

to determine which specific cognitive processes are responsible for the performance deficits in schizophrenia.

There is considerable evidence for impaired WM maintenance (Park & Holzman, 1992; Park, Püschel, Sauter, Rentsch, & Hell, 1999; Piskulic, Olver, Norman, & Maruff, 2007) and executive functions that are linked to prefrontal dysfunctions (Barch, 2005). However, performance deficits also occur with very short delays and do not necessarily increase with long delays (Gold et al., 2010; Javitt, Liederman, Cienfuegos, & Shelley, 1999; Lee & Park, 2005; Park & Holzman, 1992) pointing to a major locus of impairment at the encoding stage. Indeed, the perceptual encoding of information and its transfer into a more durable WM representation are slower (Fuller, Luck, McMahon, & Gold, 2005; Hartman, Steketee, Silva, Lanning, & McCann, 2003; Tek et al., 2002) and less precise in PSZ (Javitt, Strous, Grochowski, Ritter, & Cowan, 1997; Javitt et al., 1999; Lencz et al., 2003). These findings converge with results from electrophysiological studies (Haenschel & Linden, 2011) that implicate a role of early-stage visual processing and/or higher-level cognitive processes in abnormal encoding.

During WM encoding, the selection of task-relevant information is critical for consolidation (Sperling, 1960). The selection process allows limiting of processing to items that are salient and currently relevant in order to deal with the limited capacity of WM. Recent evidence suggests that both bottom-up cues based on perceptual stimulus features and top-down cues driven by expectations, knowledge and current goals, can increase the salience of the items to be encoded and facilitate visual WM performance in healthy participants (Fine & Minnery, 2009; Mayer, Kim, & Park, 2011; Schmidt, Vogel, Woodman, & Luck, 2002; Vogel, McCollough, & Machizawa, 2005). PSZ are also able to use simple and salient visual cues to select relevant information for WM encoding (Gold et al., 2006; Smith, Eich, Cebenoyan, & Malapani, 2011) and WM encoding improves when the task involves perceptually salient targets (Lee & Park, 2006). However, when the selection process requires a high degree of top-down control, performance is markedly impaired in PSZ both at the level of perception (Fuller et al., 2006; Gold, Fuller, Robinson, Braun, & Luck, 2007; Tanaka et al., 2007) and WM encoding (Hahn et al., 2010). Thus, difficulties in selecting relevant information or deploying attention to the relevant feature efficiently (Nestor et al., 1992; Sereno & Holzman, 1996) may result in imprecise or incorrect encoding. On the other hand, if impaired attentional processing leads to WM deficits in schizophrenia by influencing encoding, facilitating this process should improve WM performance. A recent study (Mayer et al., 2011) tested this hypothesis in healthy participants who performed a spatial delayed-response task (DRT) that manipulated orthogonally the degree of target salience and the demands on WM encoding. Target salience was manipulated by varying the degree of target novelty/familiarity (high salient = novel target, low salient = familiar target) whilst keeping the basic stimulus properties physically identical. Thus, this task was designed to probe top-down attentional processing driven by expectations rather than stimulus-dependent factors (Treisman & Gormican, 1988; Frith, 1974; Reicher, Snyder & Richards, 1976). WM load was manipulated by varying the number of targets. This study showed that increasing the salience of the targets by increasing their novelty improves spatial WM encoding in healthy participants.

We also examined different types of correct and erroneous responses. For this purpose, we asked participants to rate their level of response confidence at the end of each trial. Our classification of behavioural responses was as follows: if a subject gave the correct response with high confidence this was classified as true memory, in contrast

to a correct response that was given without confidence (correct/ not confident response). If an incorrect response was given with confidence, this was classified as false memory, in contrast to an incorrect response that was given without confidence (incorrect/ not confident response). We were particularly interested in the rate of false memory responses because false memory errors most likely reflect a problem at the encoding stage such as less precise encoding of the stimuli as demonstrated previously in PSZ (Mayer & Park, 2012). If one successfully transfers imprecisely encoded information into WM, it is possible to maintain and retrieve this information from WM, resulting in an error response that is coupled with a high degree of confidence concerning the veracity of the response. Consistent with this idea, in our previous study with healthy participants we found that the improvement in WM performance under conditions of high target salience was accompanied by a decrease in the percentage of false memory errors rather than a decrease in incorrect/ not confident responses (Mayer et al., 2011). Moreover, Lee, Folley, Gore, & Park (2008) reported similar cortical activation patterns for correct memory responses and false memory responses in a spatial DRT in PSZ, suggesting that the maintenance of the internal representation was intact whether that representation was correctly or incorrectly encoded. Therefore, if reduced WM performance in PSZ was specifically due to difficulties during encoding, we expected a higher percentage of false memory errors rather than incorrect/ not confident responses in PSZ compared to controls.

Materials and methods

Participants

Thirty-one outpatients with schizophrenia ($n = 25$) or schizoaffective disorder ($n = 6$) (PSZ) and 28 demographically matched healthy controls (HC) participated in this study. Diagnoses were made according to Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (American Psychiatric Association, 1994) using structured clinical interviews. Demographic and clinical information is summarised in Table 1. The groups were matched on age, $F(1, 58) = 1.05, p = .31$, premorbid IQ, $F(1, 58) = .79$,

Table 1. Group demographics and clinical information.

	PSZ Mean (<i>SD</i>)	HC Mean (<i>SD</i>)
Age Range	40.23 (9.10) 24–57	37.89 (8.35) 25–55
Female/male	11/20	12/16
AA : A : C : O	15 : 1 : 15 : 0	8 : 1 : 17 : 2
Handedness ^a	51.74 (61.05)	70.36 (53.33)
Years of education	13.90 (2.71)	15.29 (1.92)
IQ ^b	104.61 (9.43)	106.54 (6.84)
SAPS	14.45 (11.12)	n/a
SANS	25.06 (14.95)	n/a
BPRS	14.32 (7.28)	n/a
SPQ	n/a	10.79 (8.01)

Note: The same HC participated in the DRT and the detection task. 27 PSZ who had participated in the WM task, also performed the detection task.

^aMeasured with the Edinburgh Handedness Inventory (Oldfield, 1971).

^bMeasured with the National Adult Reading Test (Nelson, 1982). AA = African American; A = Asian; C = Caucasian; O = Other; SAPS = Scale for the Assessment of Positive Symptoms (Andreasen, 1984); SANS = Scale for the Assessment of Negative Symptoms (Andreasen, 1983); BPRS = Brief Psychiatric Rating Scale (Overall & Gorman, 1962); SPQ = Schizotypal Personality Questionnaire (Raine, 1991).

$p = .38$, and handedness, $F(1, 58) = 1.54$, $p = .22$. There was a significant difference in years of education between groups, $F(1, 58) = 5.0$, $p < .05$, which presumably reflects the effects of schizophrenia on educational attainment rather than a premorbid demographic difference.

PSZ were clinically stable (mean duration of illness: 17.57, $SD = 11.33$). Twenty-nine patients were medicated, 3 with a first-generation and 25 with a second-generation antipsychotic. Twelve patients also received antidepressants, two patients benzodiazepines, and one patient lithium. One patient was treated with antidepressant only. The mean Chlorpromazine equivalent daily dose (CPE) was 329.23 ($SD = 299.36$).

HCs were recruited from the community, had no history of DSM-IV Axis 1 disorder and were medication-free.

All subjects had normal or corrected-to-normal vision. Exclusion criteria were a history of head injury, neurological disorder or substance abuse within six months. All subjects gave written informed consent approved by the Vanderbilt University Institutional Review Board and were paid.

WM task

Stimuli, task and procedure

For details on stimuli, task and procedure see Mayer et al. (2011). Target stimuli were black “A”s of approximately $.48^\circ$ visual angle, displayed upright or upside-down on a white background (Figure 1). Stimuli were spaced evenly apart (1.9°) and appeared along an imaginary circle (4.8° radius) including 16 positions around a centrally presented fixation cross ($.36^\circ$). The positions of 0° , 90° , 180° and 270° were excluded.

The design included two within-subjects factors, target familiarity (familiar vs. unfamiliar/ novel targets) and WM load (one target vs. three targets). In the WM load 3 condition, each trial began with presenting a fixation cross at the centre position for 1s, then three targets were presented sequentially at three different positions, each for a duration of 750 ms and separated by an inter-stimulus interval of 250 ms. Within each trial, the target positions were determined pseudo-randomly with the constraint that the targets appeared in




Figure 1. Schematic diagram of the procedure and stimuli used in the WM task.

three different quadrants of the screen and that they appeared at least two positions (3.8°) farther apart from each other on the imaginary circle. In the WM load 1 condition, only the first target appeared at a position on the imaginary circle, whereas the second and the third target appeared at the centre of the screen. In the unfamiliar/ novel condition, all “A”s were presented upside-down, whereas in the familiar condition the “A”s appeared upright (Frith, 1974; Reicher et al. 1976; Shen & Reingold, 2001). After an 8-s delay interval, a question mark ($.48^\circ$ visual angle) was presented as a probe until a response was given. Participants indicated whether the position of the question mark matched one of the target positions by a left or right key press for match and non-match, respectively. Half of the trials were matches. In the non-match trials, the question mark always appeared one position further apart (1.9°) from one of the target positions along the imaginary circle to hold response difficulty constant. Participants were instructed to respond as fast and accurately as possible. Immediately after the decision, participants indicated the confidence level for their WM response by making a non-speeded response for confident and not confident. An inter-trial interval (ITI) of 3 s followed. There were 32 trials for each experimental condition presented in a randomised order and 10 practice trials.

Detection task

Participants performed a second task using the same stimuli as in the DRT, which required the detection of a target stimulus but did not place any demands on WM. The detection task allowed us to assess whether basic level perceptual processes were intact in PSZ and HC. In addition, we were interested in the effect of target novelty on visual processing and its role for efficient WM encoding. Details on the rationale for the detection task, the stimuli, task procedure and results are provided as [Supplemental Material](#).

Results

WM accuracy and RT

A repeated-measures ANOVA was performed on accuracy and RT as a function of WM load (one vs. three targets), target familiarity (familiar vs. novel target) and group (PSZ vs. HC). Response accuracy and RTs are shown in [Figure 2](#). As expected, accuracy was significantly lower in PSZ compared to HC, $F(1, 57) = 22.88, p < .001, \eta^2 = .29$. In addition, the analysis revealed a significant main effect of WM load, $F(1, 57) = 48.47, p < .001, \eta^2 = .46$, with higher accuracy for WM load 1 vs. 3. This effect was similar in both groups as reflected in the lack of a significant interaction between the factors WM load and group, $F(1, 57) = 1.13, p = .29$. The main effect of target familiarity was not significant, $F(1, 57) = .89, p = .35$. However, the interaction between target familiarity and group was significant, $F(1, 57) = 5.45, p < .05, \eta^2 = .09$, indicating differential effects of target familiarity in the two groups. In HC, there was an increase in WM accuracy for novel vs. familiar targets, $F(1, 27) = 5.31, p < .05, \eta^2 = .16$. In contrast, WM accuracy did not differ between the familiarity conditions in PSZ, $F(1, 30) = .97, p = .33$. Planned comparisons using paired *t*-tests indicated that the beneficial effect of target novelty observed in HC was significant for WM load 3, $t(27) = -2.04, p < .05$, but not for WM load 1, $t(27) = -.54, p = .59$. Interactions between WM load and target familiarity, and between all three factors were not significant (all *F*-values < 1.59 , all *p*-values $> .21$).

RTs were significantly higher in PSZ compared to HC, $F(1, 57) = 7.47, p < .01, \eta^2 = .12$. Consistent with the accuracy data, RTs increased from WM load 1 to WM load 3,




Figure 2. Mean accuracy (a) and RT (b) as a function of WM load (1 vs. 3) and target familiarity (familiar vs. novel) for healthy controls and schizophrenic patients. Vertical bars represent the standard error of the mean.

$F(1, 57) = 53.05, p < .001, \eta^2 = .48$, and this effect appeared to be stronger in PSZ than HC [WM load \times group interaction, $F(1, 57) = 5.91, p < .05, \eta^2 = .09$]. The main effect of target familiarity, $F(1, 57) = 1.90, p = .17$, and the interaction between the factors WM load and familiarity were not significant, $F(1, 57) = .40, p = .53$. Furthermore, the interaction between target familiarity and group, $F(1, 57) = 3.20, p = .08$, as well as the interaction between all three factors did not reach significance, $F(1, 57) = 2.84, p = .10$. Thus, RTs did not differ between familiarity conditions in either group indicating that the findings for accuracy were not a result of speed-accuracy trade-offs.

Confidence rating

A multifactorial repeated-measures ANOVA was conducted to test whether the percentage of type of response (true memories, correct but not confident responses, false memories, incorrect and not confident responses) differed between PSZ and CO. Specifically, we were interested whether PSZ showed an increase in the rate of false memory rather than




Figure 3. Percentage of response types across conditions for healthy controls and schizophrenic patients. Vertical bars represent the standard error of the mean.

incorrect/not confident responses compared to HC reflecting problems specifically at the stage of encoding.

As indicated by a significant interaction between the factors group and response type, $F(3, 171) = 5.53, p < .01, \eta^2 = .09$, the percentage of type of response differed between PSZ and HC. Post-hoc analyses using separate one-way ANOVAs revealed a significant group difference only in the percentage of false memories, $F(1, 58) = 18.13, p < .001$, corrected for multiple comparisons using Bonferroni correction. Across conditions, the percentage of false memory errors was about doubled in PSZ compared to HC (Figure 3). In contrast, the percentage of incorrect/not confident responses did not differ between groups, $F(1, 58) = .28, p = .60$. Overall, PSZ gave fewer true memory responses as well as fewer correct/ not confident responses than HC; however, the differences did not reach significance, $F(1, 58) = 2.67, p = .11, F(1, 58) = 1.96, p = .18$, respectively.

A subsequent analysis tested the effect of target familiarity and WM load on the percentage of false memories in PSZ and HC. The analysis was restricted to the percentage of false memories as a significant group difference emerged only for this type of response. A $2 \times 2 \times 2$ repeated-measures ANOVA revealed a significant main effect of WM load, $F(1, 57) = 11.23, p < .01, \eta^2 = .17$, with a higher percentage of false memories for load 3 than load 1 in both groups [non-significant interaction between WM load and group, $F(1, 57) = .01, p = .94$] (Figure 4). The effect of target familiarity was not significant, $F(1, 57) = 1.33, p = .25$. However, the interaction between all three factors




Figure 4. Percentage of false memory responses as a function of WM load (1 vs. 3) and target familiarity (familiar vs. novel) for healthy controls and schizophrenic patients. Vertical bars represent the standard error of the mean.

was marginally significant, $F(1, 57) = 3.09$, $p = .08$, $\eta^2 = .05$. Planned comparisons using paired t -tests indicated that this interaction effect was driven by the decrease of false memory errors for novel vs. familiar targets in the WM load 1 condition, $t(27) = 1.98$, $p = .058$, but not the WM load 3 condition, $t(27) = -.29$, $p = .78$, in HC. In contrast, in PSZ the percentage of false memories did not differ between novel and familiar targets in either the load 1, $t(27) = -.72$, $p = .48$, nor the load 3 condition, $t(27) = 1.14$, $p = .26$.

Correlations. In PSZ and HC, the percentage of false memory responses correlated negatively with overall WM performance in the low (PSZ, $r = -.83$, $p < .001$; HC, $r = -.63$, $p < .001$) and the high WM load condition (PSZ, $r = -.55$, $p < .01$; HC $r = -.66$, $p < .001$). In contrast, there was no relationship between the percentage of incorrect/ not confident responses and WM performance in PSZ (all p -values $> .12$). In HC, the percentage of incorrect/ not confident responses did not correlate with performance in the WM load 1 condition ($p = .20$), however there was a weak relationship with WM performance in the WM load 3 condition, $r = -.38$, $p < .05$.

In PSZ, there was no relationship between response accuracy and symptom severity (all p -values $> .14$). However, the increase in false memories from WM load 1 to load 3 significantly correlated with the severity of negative symptoms, $r = .48$, $p < .01$.

There was a trend for a negative correlation between response accuracy and the daily CPE dose only in the condition WM load 1/novel targets, $r = -.38$, $p = .06$. In all other experimental conditions, we did not find significant correlations between task performance and medication doses (all p -values $> .11$). There was also no relationship between WM accuracy and duration of illness (all p -values $> .11$).

Discussion

In this study, we investigated the effect of target salience on spatial WM in schizophrenia by manipulating the novelty of the targets to be encoded in a DRT. WM performance was markedly impaired in PSZ. Increasing target salience by increasing the novelty improved WM performance in HC but not in PSZ. Thus, in contrast to our hypothesis, PSZ failed to benefit from increased novelty of the target during WM encoding. In addition, the beneficial effect of target novelty on visual processing as assessed in the detection task was positively correlated with performance in the DRT in HC rather than PSZ. Thus, healthy participants who benefit more from novel targets in the detection task showed better WM performance (see [Supplemental Material](#), Figure S1a).

The findings from the confidence rating revealed that impaired WM performance in PSZ was mainly due to an increase in the number of false memory responses. The percentage of false memories increased with WM load and was doubled in PSZ compared to HC, whereas the amount of incorrect responses that were given without being confident did not differ between groups. Also, the percentage of false memory responses rather than the percentage of incorrect/ not confident responses correlated negatively with WM performance in both WM load conditions. This finding indicates that the degree to which participants made false memory errors was related to their degree of WM reduction. In HC, the percentage of false memory errors was reduced for novel vs. familiar targets; however, this effect was small and appeared only in the load 1 condition. More importantly, the percentage of false memory responses also correlated negatively with the difference in RT between novel and familiar targets in the detection task when calculated across participants (Figure S1b). Thus, those participants who benefitted little from novel targets in the detection task made more false memory errors in the WM task.

This provides some evidence that the increased rate of false memory errors likely reflects difficulties in deploying attention, which then may lead to inefficient WM encoding. This is consistent with our previous findings indicating that the impairments in spatial WM observed in PSZ as well as healthy first-degree relatives of PSZ can be attributed at least to some degree, to deficits in processes associated with WM encoding such as less precise encoding (Lee & Park, 2005; Mayer & Park, 2012). This finding is also consistent with the results from neuroimaging studies that compared true memory and false memory responses in PSZ and observed similar activation patterns in the prefrontal cortex for both types suggesting that the mechanisms that support the maintenance of the internal representation were intact whether that representation was correctly or incorrectly encoded (Lee et al., 2008).

Taken together, the present findings support and extend previous reports on the relevance of processes associated with the early phase of encoding for WM deficits in schizophrenia (Fuller et al., 2005; Haenschel & Linden, 2011; Hahn et al., 2010; Hartman et al., 2003; Javitt et al., 1999; Lee & Park, 2005; Lencz et al., 2003; Mayer, Fukuda, Vogel, & Park, 2012; Mayer & Park, 2012; Tek et al., 2002). Analysing the overall performance and types of responses we demonstrate that dysfunctions of visual attention in the service of WM encoding, rather than a failure of WM storage per se, contribute to WM impairments in schizophrenia.

It might be argued that the deficit in WM performance in PSZ was due to slowed perceptual encoding rather than attentional processing. However, given that the target detection time in PSZ was well below the target exposure time of 750 ms (see [Supplemental Material](#)), we can rule out that slowed perceptual encoding was a limiting factor for WM performance in our task.

The increased rate of false memories might reflect an overall response bias for confident responses in PSZ rather than a specific deficit of attentional processing. In this case, we would have expected a similar distribution of confident and not confident responses among correct and incorrect responses in PSZ. The results were not consistent with this hypothesis. Across conditions, 84% of the correct responses were true memories and 16% of the correct responses were given without being confident. In contrast, among the errors about 75% were false memory errors, whereas 25% were incorrect and not confident responses.

Our findings pinpoint a circumscribed deficit in the processing of novelty in PSZ. It is important to note that in the present task the influence of low-level physical features on target salience was minimised as novel and familiar targets differed only with regard to a local change in orientation. The attentional salience of the novel target was likely driven by expectations regarding a previously stored concept of the letter “A” (Frith, 1974; Mayer et al., 2011; Reicher et al., 1976; Shen & Reingold, 2001). Thus, the patients’ deficit in assigning attentional salience to novel targets in order to facilitate WM encoding adds to the growing body of evidence that top-down processes function abnormally in schizophrenia when visual inputs are selected for further perceptual (Fuller et al., 2006; Gold et al., 2007; Tanaka et al., 2007) and memory processing (Hahn et al., 2010). In contrast, when the selection process is driven by perceptually salient stimulus features and/or the distractors do not strongly compete for attentional resources, attentional selection is intact in PSZ and facilitates WM encoding (Gold et al., 2006; Lee & Park, 2006; Smith et al., 2011).

All PSZ except for three were taking antipsychotic medication. However, the profile of WM deficit observed in PSZ cannot be explained solely in terms of medication. First,

we did not find a consistent relationship between WM performance and daily medication dosage. Second, preliminary data in a group of healthy relatives (REL) of PSZ who were medication-free showed that WM accuracy was also lower in this group compared to HC and similar to PSZ this deficit was also accompanied by an increase in false memories (see [Supplemental Material](#)). Consistent with previous reports (Mayer & Park, 2012) these findings suggest that PSZ and REL might be comparable in terms of spatial WM deficits as well as the underlying mechanisms.

Understanding the processes that contribute to impaired WM in schizophrenia is crucial in the search for cognitive remediation strategies. The present findings suggest that non-mnemonic attentional processes in the service of WM encoding represent an important target for improving WM.

Acknowledgements

This work was supported by the National Institutes of Health [grant number R01 MH073028 to S.P. and grant number P30 HD15052 to the Vanderbilt Kennedy Center for Research on Human Development]. We thank Lindsey Gilling McIntosh for help with data collection, Heath Nichols, Katy Thakkar, and Joel Peterman for their assistance with clinical interviews and subject recruitment, and Professor Stephan Heckers with his help in recruitment.

Supplementary material

Supplementary (Figure SI and S2 and further content) is available via the 'Supplementary' tab on the article's online page (<http://dx.doi.org/10.1080/13546805.2013.854199R>).

References

- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Andreasen, N. C. (1983). *The scale for the assessment of negative symptoms (SANS)*. Iowa City, IA: The University of Iowa.
- Andreasen, N. C. (1984). *The scale for the assessment of positive symptoms (SAPS)*. Iowa City, IA: The University of Iowa.
- Barch, D. M. (2005). The cognitive neuroscience of schizophrenia. *Annual Review of Clinical Psychology, 1*, 321–353. doi:10.1146/annurev.clinpsy.1.102803.143959
- Bledowski, C., Kaiser, J., & Rahm, B. (2010). Basic operations in working memory: contributions from functional imaging studies. *Behavioural Brain Research, 214*, 172–179. doi:10.1016/j.bbr.2010.05.041
- Cervellione, K. L., Burdick, K. E., Cottone, J. G., Rhinewine, J. P., & Kumra, S. (2007). Neurocognitive deficits in adolescents with schizophrenia: Longitudinal stability and predictive utility for short-term functional outcome. *Journal of the American Academy of Child and Adolescent Psychiatry, 46*, 867–878. doi:10.1097/chi.0b013e318054678d
- Fine, M. S., & Minnery, B. S. (2009). Visual salience affects performance in a working memory task. *Journal of Neuroscience, 29*, 8016–8021. doi:10.1523/JNEUROSCI.5503-08.2009
- Frith, U. (1974). A curious effect with reversed letters explained by a theory of schema. *Attention & Psychophysics, 16*, 113–116. doi:10.3758/BF03203262
- Fuller, R. L., Luck, S. J., Braun, E. L., Robinson, B. M., McMahon, R. P., & Gold, J. M. (2006). Impaired control of visual attention in schizophrenia. *Journal of Abnormal Psychology, 115*, 266–275. doi:10.1037/0021-843X.115.2.266
- Fuller, R. L., Luck, S. J., McMahon, R. P., & Gold, J. M. (2005). Working memory consolidation is abnormally slow in schizophrenia. *Journal of Abnormal Psychology, 114*, 279–290. doi:10.1037/0021-843X.114.2.279
- Glahn, D. C., Therman, S., & Manninen, M. (2003). Spatial working memory as an endophenotype for schizophrenia. *Biological Psychiatry, 53*, 624–626. doi:10.1016/S0006-3223(03)01641-4

- Gold, J. M., Fuller, R. L., Robinson, B. M., Braun, E. L., & Luck, S. J. (2007). Impaired top-down control of visual search in schizophrenia. *Schizophrenia Research*, *94*, 148–155. doi:[10.1016/j.schres.2007.04.023](https://doi.org/10.1016/j.schres.2007.04.023)
- Gold, J. M., Fuller, R. L., Robinson, B. M., McMahon, R. P., Braun, E. L., & Luck, S. J. (2006). Intact attentional control of working memory encoding in schizophrenia. *Journal of Abnormal Psychology*, *115*, 658–673. doi:[10.1037/0021-843X.115.4.658](https://doi.org/10.1037/0021-843X.115.4.658)
- Gold, J. M., Hahn, B., Zhang, W. W., Robinson, B. M., Kappenman, E. S., Beck, V. M., & Luck, S. J. (2010). Reduced capacity but spared precision and maintenance of working memory representations in schizophrenia. *Archives of General Psychiatry*, *67*, 570–577. doi:[10.1001/archgenpsychiatry.2010.65](https://doi.org/10.1001/archgenpsychiatry.2010.65)
- Goldman-Rakic, P. S. (1994). Working memory dysfunction in schizophrenia. *Journal of Neuropsychiatry and Clinical Neurosciences*, *6*, 348–357.
- Haenschel, C., & Linden, D. (2011). Exploring intermediate phenotypes with EEG: working memory dysfunction in schizophrenia. *Behavioural Brain Research*, *216*, 481–495. doi:[10.1016/j.bbr.2010.08.045](https://doi.org/10.1016/j.bbr.2010.08.045)
- Hahn, B., Robinson, B. M., Kaiser, S. T., Harvey, A. N., Beck, V. M., Leonard, C. J., & Gold, J.M. (2010). Failure of schizophrenia patients to overcome salient distractors during working memory encoding. *Biological Psychiatry*, *68*, 603–609. doi:[10.1016/j.biopsych.2010.04.014](https://doi.org/10.1016/j.biopsych.2010.04.014)
- Hartman, M., Steketee, M., Silva, S., Lanning, K., & McCann, H. (2003). Working memory and schizophrenia: evidence for slowed encoding. *Schizophrenia Research*, *59*, 99–113. doi:[10.1016/S0920-9964\(01\)00366-8](https://doi.org/10.1016/S0920-9964(01)00366-8)
- Javitt, D. C., Liederman, E., Cienfuegos, A., & Shelley, A.-M. (1999). Panmodal processing imprecision as a basis for dysfunction of transient memory storage systems in schizophrenia. *Schizophrenia Bulletin*, *25*, 763–775. doi:[10.1093/oxfordjournals.schbul.a033417](https://doi.org/10.1093/oxfordjournals.schbul.a033417)
- Javitt, D. C., Strous, R. D., Grochowski, S., Ritter, W., & Cowan, N. (1997). Impaired precision, but normal retention, of auditory sensory (“echoic”) memory information in schizophrenia. *Journal of Abnormal Psychology*, *106*, 315–324. doi:[10.1037/0021-843X.106.2.315](https://doi.org/10.1037/0021-843X.106.2.315)
- Lee, J., Folley, B. S., Gore, J., & Park, S. (2008). Origins of spatial working memory deficits in schizophrenia: An event-related fMRI and near-infrared spectroscopy study. *PLoS One*, *3*, e1760. doi:[10.1371/journal.pone.0001760](https://doi.org/10.1371/journal.pone.0001760)
- Lee, J., & Park, S. (2005). Working memory impairments in schizophrenia: A meta-analysis. *Journal of Abnormal Psychology*, *114*, 599–611. doi:[10.1037/0021-843X.114.4.599](https://doi.org/10.1037/0021-843X.114.4.599)
- Lee, J., & Park, S. (2006). The role of stimulus salience in CPT-AX performance of schizophrenia patients. *Schizophrenia Research*, *81*, 191–197. doi:[10.1016/j.schres.2005.08.015](https://doi.org/10.1016/j.schres.2005.08.015)
- Lenz, T., Bilder, R. M., Turkel, E., Goldman, R. S., Robinson, D., Kane, J. M., & Lieberman, J. A. (2003). Impairments in perceptual competency and maintenance on a visual delayed match-to-sample test in first-episode schizophrenia. *Archives of General Psychiatry*, *60*, 238–243. doi:[10.1001/archpsyc.60.3.238](https://doi.org/10.1001/archpsyc.60.3.238)
- Mayer, J. S., Fukuda, K., Vogel, E. K., & Park, S. (2012). Impaired contingent attentional capture predicts reduced working memory capacity in schizophrenia. *PLoS One*, *7*, e48586. doi:[10.1371/journal.pone.0048586](https://doi.org/10.1371/journal.pone.0048586)
- Mayer, J. S., Kim, J., & Park, S. (2011). Enhancing visual working memory encoding: The role of target novelty. *Visual Cognition*, *19*, 863–885. doi:[10.1080/13506285.2011.594459](https://doi.org/10.1080/13506285.2011.594459)
- Mayer, J. S., & Park, S. (2012). Working memory encoding and false memory in schizophrenia and bipolar disorder in a spatial delayed response task. *Journal of Abnormal Psychology*, *121*, 784–794. doi:[10.1037/a0028836](https://doi.org/10.1037/a0028836)
- Meyer-Lindenberg, A., & Weinberger, D. R. (2006). Intermediate phenotypes and genetic mechanisms of psychiatric disorders. *Nature Reviews Neuroscience*, *7*, 818–827. doi:[10.1038/nrn1993](https://doi.org/10.1038/nrn1993)
- Mitropoulou, V., Harvey, P. D., Zegarelli, G., New, A. S., Silverman, J. M., & Siever, L. J. (2005). Neuropsychological performance in schizotypal personality disorder: importance of working memory. *American Journal of Psychiatry*, *162*, 1896–1903. doi:[10.1176/appi.ajp.162.10.1896](https://doi.org/10.1176/appi.ajp.162.10.1896)
- Myles-Worsley, M., & Park, S. (2002). Spatial working memory deficits in schizophrenia patients and their first degree relatives from Palau, Micronesia. *American Journal of Medical Genetics*, *114*, 609–615. doi:[10.1002/ajmg.10644](https://doi.org/10.1002/ajmg.10644)
- Nelson, H. E. (1982). *National adult reading test*. Windsor: NFER-Nelson.

- Nestor, P., Faux, S., McCarley, R., Penhune, V., Shenton, M., Pollak, S., & Sands, S. F. (1992). Attentional cues in chronic schizophrenia: Abnormal disengagement of attention. *Journal of Abnormal Psychology, 101*, 682–689. doi:10.1037/0021-843X.101.4.682
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia, 9*, 97–113. doi:10.1016/0028-3932(71)90067-4
- Overall, J. E., & Gorman, D. R. (1962). The brief psychiatric rating scale. *Psychological Reports, 10*, 799–812. doi:10.2466/pr0.1962.10.3.799
- Park, S., & Holzman, P. S. (1992). Schizophrenics show spatial working memory deficits. *Archives of General Psychiatry, 49*, 975–982. doi:10.1001/archpsyc.1992.01820120063009
- Park, S., Holzman, P. S., & Goldman-Rakic, P. S. (1995). Spatial working memory deficits in the relatives of schizophrenic patients. *Archives of General Psychiatry, 52*, 821–828. doi:10.1001/archpsyc.1995.03950220031007
- Park, S., Püschel, J., Sauter, B., Rentsch, M., & Hell, D. (1999). Spatial working memory deficits and clinical symptoms in schizophrenia: A 4-month follow-up study. *Biological Psychiatry, 46*, 392–400. doi:10.1016/S0006-3223(98)00370-9
- Piskulic, D., Olver, J. S., Norman, T. R., & Maruff, P. (2007). Behavioural studies of spatial working memory dysfunction in schizophrenia: A quantitative literature review. *Psychiatry Research, 150*, 111–121. doi:10.1016/j.psychres.2006.03.018
- Raine, A. (1991). The SPQ: A scale for the assessment of schizotypal personality based on DSMIII-R criteria. *Schizophrenia Bulletin, 17*, 555–564. doi:10.1093/schbul/17.4.555
- Reicher, G. M., Snyder, C. R., & Richards, J. T. (1976). Familiarity of background characters in visual scanning. *Journal of Experimental Psychology: Human Perception and Performance, 2*, 522–530. doi:10.1037/0096-1523.2.4.522
- Saperstein, A. M., Fuller, R. L., Avila, M. T., Adami, H., McMahon, R. P., Thaker, G. K., & Gold, J. M. (2006). Spatial working memory as a cognitive endophenotype of schizophrenia: Assessing risk for pathophysiological dysfunction. *Schizophrenia Bulletin, 32*, 498–506. doi:10.1093/schbul/sbj072
- Schmidt, B. K., Vogel, E. K., Woodman, G. F., & Luck, S. J. (2002). Voluntary and automatic attentional control of visual working memory. *Perception & Psychophysics, 64*, 754–763. doi:10.3758/BF03194742
- Sereno, A. B., & Holzman, P. S. (1996). Spatial selective attention in schizophrenic, affective disorder, and normal subjects. *Schizophrenia Research, 20*, 33–50. doi:10.1016/0920-9964(95)00077-1
- Shen, J., & Reingold, E. M. (2001). Visual search asymmetry: The influence of stimulus familiarity and low-level features. *Perception & Psychophysics, 63*, 464–475. doi:10.3758/BF03194413
- Smith E. E., Eich, T. S., Cebenoyan, D., & Malapani, C. (2011). Intact and impaired cognitive-control processes in schizophrenia. *Schizophrenia Research, 126*, 132–137. doi:10.1016/j.schres.2010.11.022
- Smith, C. W., Park, S., & Cornblatt, B. (2006). Spatial working memory deficits in adolescents at clinical high risk for schizophrenia. *Schizophrenia Research, 81*, 211–215. doi:10.1016/j.schres.2005.09.019
- Snitz B. E., MacDonald, A. W. 3rd, & Carter, C. S. (2006). Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: A meta-analytic review of putative endophenotypes. *Schizophrenia Bulletin, 32*, 179–194. doi:10.1093/schbul/sbi048
- Sperling G. (1960). The information available in brief visual presentations. *Psychological Monographs: General and Applied, 74*, 1–29. doi:10.1037/h0093759
- Tanaka, G., Mori, S., Inadomi, H., Hamada, Y., Ohta, Y., & Ozawa, H., (2007). Clear distinction between preattentive and attentive process in schizophrenia by visual search performance. *Psychiatry Research, 149*, 25–31. doi:10.1016/j.psychres.2006.01.014
- Tek, C., Gold, J., Blaxton, T., Wilk, C., McMahon, R. P., & Buchanan, R. W. (2002). Visual perceptual and working memory impairments in schizophrenia. *Archives of General Psychiatry, 59*, 146–153. doi:10.1001/archpsyc.59.2.146
- Treisman, A., & Gormican, S. (1988). Feature analysis in early vision: Evidence from search asymmetries. *Psychological Review, 1*, 15–48. doi:10.1037/0033-295X.95.1.15
- Vogel, E. K., McCollough, A. W., & Machizawa, M. G. (2005). Neural measures reveal individual differences in controlling access to working memory. *Nature, 438*, 500–503. doi:10.1038/nature04171